



5-7-2018

# Kinetic studies on the aqueous chemistry of the glutathione-methyleneoxindole conjugate to model its inhibition of glyoxalase I

Ariane Borges

Follow this and additional works at: [http://vc.bridgew.edu/honors\\_proj](http://vc.bridgew.edu/honors_proj)

 Part of the [Chemistry Commons](#)

## Recommended Citation

Borges, Ariane. (2018). Kinetic studies on the aqueous chemistry of the glutathione-methyleneoxindole conjugate to model its inhibition of glyoxalase I. In *BSU Honors Program Theses and Projects*. Item 300. Available at: [http://vc.bridgew.edu/honors\\_proj/300](http://vc.bridgew.edu/honors_proj/300)  
Copyright © 2018 Ariane Borges

Kinetic studies on the aqueous chemistry of the glutathione-methyleneoxindole conjugate  
to model its inhibition of glyoxalase I

Ariane Borges

Submitted in Partial Completion of the  
Requirements for Departmental Honors in Chemistry

Bridgewater State University

May 7, 2018

Dr. Edward Brush, Thesis Advisor  
Dr. Saritha Nellutla, Committee Member  
Dr. Samer Lone, Committee Member

**Kinetic studies on the aqueous chemistry  
of the glutathione-methyleneoxindole  
conjugate to model its inhibition of  
glyoxalase I**

*Ariane Borges*

Undergraduate Honors Thesis

Mentor:

Dr. Edward J. Brush

Bridgewater State University  
Department of Chemical Sciences  
Bridgewater, MA 02325

This page intentionally left blank.

## Abstract

Glyoxalase I (GxI) is an enzyme that is part of the Glyoxalase system which is responsible for the conversion of methylglyoxal (MG), a byproduct of glycolysis, to lactic acid using glutathione (GSH) as a co-substrate. Methylglyoxal and GSH come together to form the hemithioacetal substrate, which is then transformed into (S)-D-lactoylglutathione. Since MG is a highly reactive compound known to induce cell apoptosis, increasing MG levels in cells by inhibiting GxI should increase cell death. Research into the design, synthesis and testing of different inhibitors of GxI support this hypothesis. We have designed and synthesized a potential inhibitor of GxI, the glutathione (GSH)-3-methyleneoxindole (MOI) conjugate (GSMOI). Upon binding to the active site of GxI, we propose that GxI will catalyze a proton transfer elimination reaction on GSMOI, releasing GSH and producing MOI which will irreversibly alkylate and inhibit GxI.

To test this hypothesis, the purpose of this research was to determine the chemical mechanism by which GSMOI might undergo elimination (E1, E2 or E1cb), and how this reaction might be controlled and affected under physiological conditions. MOI and GSMOI have been synthesized, and their purity and structure characterized using UV-Vis spectroscopy and Nuclear Magnetic Resonance (NMR) spectrometry. The chemical kinetics of the elimination reaction of GSH from GSMOI were studied by monitoring the formation of either GSH or MOI by UV-Vis spectroscopy. The E1 elimination is a two-step mechanism where changes in pH should not affect the rate of the reaction, while in E2 and E1cb mechanisms the rate was expected to increase with increase in pH. Incubation of GSMOI at 25°C in phosphate buffers of different pH results in a significant drop in reaction rate below pH 6. This suggests that changes in pH affect the rate of reaction, and that the elimination is following either E2 or E1cb mechanism. To distinguish between these mechanisms, we used <sup>1</sup>H NMR to study GSMOI proton exchange in D<sub>2</sub>O solvent. These results conclusively showed that the C-3 proton of GSMOI rapidly exchanged with D<sub>2</sub>O solvent, without the corresponding formation of MOI, leading to the conclusion that the reaction is following an E1cb conjugate base intermediate mechanism. This research was supported by the Department of Chemical Sciences at Bridgewater State University and by a summer 2017 grant from the BSU Adrian Tinsley Program.

## **Preface**

Every research project starts with some unsolved question, or some type of mystery in a specific subject or topic. This thesis, presented to the Department of Chemical Sciences in partial fulfillment of the Bachelor of Science degree with Honors in Chemistry (Biochemistry concentration), is a study into the mystery of the aqueous chemistry of the glutathione-methyleneoxindole conjugate in order to understand its solution chemistry, and to model its physiological reactions as a potential inhibitor of the anti-cancer target enzyme, Glyoxalase I.

The introduction describes background about the Glyoxalase pathway and how it uses glutathione to metabolize its cytotoxic substrate, methylglyoxal. The possible role of the Glyoxalase system in the control of cell growth is also discussed, and how Glyoxalase inhibitors have been proposed to increase methylglyoxal levels and induce apoptosis in cancer cells. That leads to our proposed mechanism-based Glyoxalase inhibitor, the Glutathione-Methyleneoxindole conjugate (GSMOI). We have proposed that GSMOI activation by a Glyoxalase-catalyzed elimination reaction could lead to the formation of the alkylating agent, 3-methyleneoxindole. Understanding the aqueous chemistry and the mechanism of GSMOI elimination reactions is crucial to understanding its potential role in Glyoxalase I inhibition. The remaining sections, Methodology, Results, Discussion and Conclusions, describe the research conducted with Dr. Brush at Bridgewater State University. This research presents experimental evidence for an E1cb elimination mechanism for GSMOI, which leads to a discussion on the relevance of this new finding to the potential activation of GSMOI and its inhibition of Glyoxalase I. Based on the work described in this thesis, additional questions have been proposed inviting new researchers to this project.

So, I hope that this thesis serves as a guide to all of those reading it and new researchers in general.

### **Thesis Committee Approval (as to style and content):**

---

**Edward J. Brush, Ph.D., Department of Chemical Sciences**

---

**Saritha Nellutla, Ph.D., Department of Chemical Sciences**

---

**Samer Lone, Ph.D., Department of Chemical Sciences**

## Acknowledgements

First, I would like to thank my mentor, Dr. Edward J. Brush for all his guidance and support throughout my research, and for mentoring me. I would like to thank him for making me aware of great opportunities and for always being patient and kind with me. Without him I wouldn't have discovered my passion for chemistry or research and I wouldn't have written this thesis. I am eternally grateful for everything he has done for me.

I would also like to thank my adviser for my Biology degree Dr. Jeffery Bowen, for always being helpful and guiding me through the burden of picking the right classes at the right times. I would like to thank all the professors that made a difference in my undergraduate journey both in the humanities and in the sciences. Each one of them pushed me to become a better version of myself and a better student. I would like to thank them all for their constructive criticism, life lessons and for always being there for me.

I would like to thank my colleagues in the Biology department who have graduated, for being the best upper classmates anyone could wish for. I have a profound thanks to my chemistry colleagues for always being supportive and helpful. I am very glad I got to meet each and every one of you and my college journey wouldn't have been the same without you.

Lastly, I would like to thank all my family and friends, especially my parents for always being very supportive, I wouldn't be able to accomplish this if it weren't for your unconditional support and guidance.

The following research was supported by the BSU Adrian Tinsley Program. The JEOL ECX-400 MHz NMR was obtained through NSF-MRI grant 0421081. Thesis is being submitted in partial fulfillment of a Bachelor of Science degree with Departmental Honors in Chemistry (Biochemistry concentration).

## Table of Contents

Abstract.....	3
1. Introduction	
1.1. Glyoxalase System.....	7-11
1.2. Inhibition of Glyoxalase I.....	12
1.3. Glutathione Methyleneoxindole conjugation (GSMOI).....	13-17
1.4. Significance of this Work.....	17
2. Materials and Methods	
2.1. Synthesis of 3-Bromooxindol-3-acetic acid.....	18-19
2.2. Synthesis of 3-methyleoxindole (MOI) from 3-Bromooxindole-3-acetic acid (BOAA).....	19-20
2.3. Synthesis of Glutathione-3-Methyleneoxindole Conjugate, GSMOI from 3-methyleoxindole.....	20-21
2.4. Characterization using NMR.....	21
2.5. UV-Vis Kinetic Studies.....	21-22
2.6. Determination of reaction order.....	22
2.7. Effect of pH on the rate of GSH elimination .....	23
2.8. Effect of potential catalyst .....	23
2.9. Effect of buffer concentration on rate constant K.....	23
2.10. GSMOI C-3 Proton exchange kinetics.....	24
3. Results and discussion.....	25-36
4. Conclusion.....	37-41
5. Future Work.....	42
6. Bibliography.....	43-44




# 1. Introduction

## 1.1 Glyoxalase System

### 1.1.1 Discovery and first insight


In 1913 researchers discovered the formation of lactic acid from methylglyoxal. At the time, they believed this conversion to be associated with one enzyme which they called Glyoxalase. However, in 1951 new studies revealed the need for two enzymes and a co-factor for the conversion of methylglyoxal to lactic acid. The Glyoxalase/Glycolysis pathway is shown in Figure 1. The first enzyme in the Glyoxalase pathway is Glyoxalase I (GxI or GLO I) which is responsible for the conversion of methylglyoxal into (S)-D-lactoylglutathione. Then Glyoxalase II (GxII) converts (S)-D-lactoylglutathione into D-lactate, which can then be converted into pyruvate. The glycolysis pathway uses pyruvate as the main substrate to generate ATP through aerobic respiration. The glyoxalase pathway is found in all cells and has two main functions: as an antioxidant defense and in the production of ATP through an alternate energy source.



**Figure 1:** The reaction schematic for the glyoxalase enzymatic pathway.

### ***1.1.2 Methylglyoxal: Biosynthesis, damage and detoxification***


Methylglyoxal (MG) is one of the byproducts of oxidative metabolism in cells and since it is a highly reactive compound it is detoxified by the glyoxalase system present in cells, as described in Figures 2 and 3. Methylglyoxal is highly reactive and is inevitably formed as a by-product of glycolysis. MG can be produced by enzymatic and non-enzymatic methods, and its rate of formation is dependent on the organism, tissue and cell type. In glycolysis, MG is produced by the fragmentation of two triosephosphates, specifically glyceraldehyde-3-phosphate (GAP) and dihydroxyacetone phosphate (DHAP), Figure 2. This is the main route of generating MG, but it can also be generated by metabolism of protein and fatty acids. Studies show that about 0.1-0.4% of the glycolytic flux results in MG production (Kalapos, 2008). MG is very reactive, and about 90–99% of cellular MG is found bound to biomacromolecules. For example, Kalapos demonstrated that when 1  $\mu\text{M}$  of MG was added to human plasma it bound irreversibly to the plasma protein within 24 h of incubation at 37°C. The concentration of MG in cells, both free and bound to macromolecules, can be up to 300 $\mu\text{M}$  implying that MG has a stronger potency at the site of production and may have little to no potency at distant sites. As shown in Figure 3, Methylglyoxal is thought to be involved in mediating the diffusion of advanced glycation end-products (AGEs) through the cell membrane. These AGEs are associated with different pathologies such as Parkinson's disease, Alzheimer's disease, and diabetes. MG is also known to react with lipids, nucleic acids and some residues of proteins involved in the formation of AGEs (Okado 1996).



**Figure 2.** The enzymatic pathway comprised of Glyoxalase I and Glyoxalase II responsible for the conversion of methylglyoxal to lactic acid, and Glycolysis Pathway. Glucose Metabolism: Glycation and Methylation, <https://www.realizehealth.com.au/2015/08/18/glucose-metabolism-glycation-and-methylation/> (Accessed March 31, 2018)

Under normal circumstances, cells are protected against MG toxicity by different enzymatic pathways such as glyoxalase, aldose reductase, aldehyde dehydrogenase and carbonyl reductase pathways. However, the glyoxalase pathway is the most effective pathway (Thornalley, 1993; Kalapos, 1999; Vander Jagt and Hunsaker, 2003). Evidence has also been gathered to support the idea that MG promotes the activation of enzymes involved in apoptosis such as protein kinase C, c-Jun N-terminal Kinases and p-38 mitogen-activated protein kinase add reference. A recent study showed that in apoptosis mediated by reactive oxygen species MG can induce apoptosis in a concentration-dependent manner. All the data gathered was specific to bovine retinal

pericyte and therefore more studies involving different cell types from a variety of organisms need to be done in order to make a broader conclusion. In addition, it was concluded that high levels of MG can cause apoptosis through an oxidative stress mechanism and that the nuclear activation of the transcription factor nuclear factor  $\kappa$ B (NF- $\kappa$ B) is involved in the apoptosis process (Okado 1996). Other evidence has shown that MG prevents cancer cells from growing by inhibiting mitochondrial respiration (Okado 1996, Wu 2002).




**Figure 3.** The mechanism of biosynthesis and detoxification of methylglyoxal and the damage caused by methylglyoxal. Coordinated Actions of Glyoxalase and Antioxidant Defense Systems in Conferring Abiotic Stress Tolerance in Plants. <http://www.mdpi.com/1422-0067/18/1/200/htm> (Accessed May 8, 2018)

### 1.1.3 Structure of Glyoxalase I

Glyoxalase I (GxI) is the first enzyme in the glyoxalase pathway, and is thought to regulate cell growth and proliferation by controlling the amount of MG in living cells. This system is present in the cytosol of cells where GxI catalyzes the formation of (S)-D-lactoylglutathione from

the hemithioacetal which is formed in a non-enzymatic reaction between MG and GSH (see Figure 2). Then, GxII converts (S)-D-lactoylglutathione to lactic acid which is relatively non-toxic and regenerates the co-substrate, GSH. The glyoxalase pathway is highly conserved throughout biological life and is present also in cell organelles and mitochondria. This system is active from embryogenesis until cell death, and plays an important role in cell growth and division.

Methylglyoxal, glyoxal and other acyclic  $\alpha$ -oxoaldehydes are the main substrates of GxI. There is a slight variation in the structure of GxI between organisms. Human GxI is a dimeric  $Zn^{2+}$  metalloenzyme whereas *Escherichia coli* GxI is a  $Ni^{2+}$  metalloenzyme. Both of these metals are essential for catalysis. Each monomer consists of two structurally equivalent domains and the active site is situated at the dimer interface. Human GxI, shown in the ribbon structure in Figure 4, has a molecular mass of 42kDa and is regulated by phosphorylation in four different sites.



**Figure 4.** Crystal structure of Human Glyoxalase I, molecular mass of 42KDa. Image generated using FirstGlance software.


## ***1.2 Inhibition of Glyoxalase I***

### ***1.2.1 Hallmarks of Apoptosis***

GxI has been extensively studied in the past years because its inhibitors are shown to be potential cancer therapeutic agents. By inhibiting GxI, MG levels should increase, leading to an increase in cell death by apoptosis or necrosis. The cells of a multicellular organism form an organized community, and this community is extremely regulated. The regulation involves controlling the rates of cell division and cell death (White, 2004). When a cell is no longer viable and is not contributing to the overall well-being of the community, this cell undergoes apoptosis, which is a process of programmed cell death that occurs under normal physiological conditions. This form of cell death is more organized when compared to necrosis, which occurs when the cell encounters an extreme variance in physiological conditions. Both processes of cell death are part of the normal growth and development of an individual, and many pathologies can arise if these processes are inhibited (White 2004, Osborne 1996). Hence, if GxI is inhibited, the level of MG will increase, and therefore the rate of cell death by apoptosis will increase. Different competitive GxI inhibitors, designed based on the active site structure, have been synthesized to test this hypothesis. These inhibitors have been found to have anti-proliferative activity and suggest that the increased levels of MG do indeed prevent tumor growth. The best competitive inhibitors of GxI were found to be GSH derivatives of small, hydrophobic compounds.

### 1.3 Glutathione-Methyleneoxindole conjugate (GSMOI), a potential inhibitor of GxI


The Glutathione-Methyleneoxindole conjugate (GSMOI), is a molecule formed by conjugating glutathione (GSH) and 3-methyleneoxindole (MOI) as shown in Figure 5. MOI is proposed to be the actual enzyme inhibitor due to nucleophile alkylation. However, it cannot be used directly as a therapeutic agent because it is chemically reactive and it will most likely alkylate and inactivate other enzymes and biomolecules that are not part of the glyoxalase system, thereby disrupting other physiological processes. In addition, MOI is a nonpolar molecule that is not soluble in water and therefore is not soluble in the bloodstream. Previous preliminary studies indicated that GSMOI is both a competitive and mechanism-based inhibitor of GxI, however more in-depth work is needed to shed light on these findings. (Ivancic and Brush, unpublished results).



**Figure 5:** Synthesis of GSMOI from glutathione and 3-methyleneoxindole.

### 1.3.1 Synthesis of GSMOI

As mentioned above, MOI is not a good choice as a therapeutic agent because of its poor aqueous solubility and non-selectivity as an inhibitor and alkylating agent. By conjugating MOI with GSH, this will make it more specific to target GxI and inhibit it via a mechanism-based inactivation. Once bound to the active site of GxI, it is hypothesized that GxI will catalyze the elimination of GSH from GSMOI, and forming MOI. As MOI is a potent alkylating agent for reactive thiols, the free MOI has high chance to selectively alkylate a cysteine residue at the active site via Michael addition, Figure 6. As can be seen from Figure 6, the proposed formation of GSH and MOI from GSMOI by GxI is an example of elimination reaction because of the characteristic features such as the presence of a leaving group, loss of a proton and the formation of a product with a double bond (MOI). Understanding how this reaction occurs, and what type of elimination reaction it is, will be beneficial for predicting products of similar reactions, and it may provide information to understand how the reaction can be controlled or affected.



**Figure 6:** Proposed inactivation of GxI by GSMOI through GSH elimination and active site cysteine alkylation by MOI.

### 1.3.2 Elimination of MOI from GSMOI




The purpose of this thesis project was to investigate and better understand the mechanism by which GSH is eliminated from GSMOI, producing free MOI. Understanding this mechanism may require studies on how the reaction rate might be affected by factors such as pH, temperature



and potential acid or base catalysts. This work depended on the ability to develop an assay to quantitatively study the rate of the elimination reaction, under conditions that would allow the study of factors that affect the rate limiting step. This turned out to be a challenge that was solved. The interpretation of rate data required understanding of the chemical basis behind the three most common biological elimination mechanisms that could explain the elimination of GSH and formation of MOI: E1, E2 or E1cb (cf. Figure 7). (Smith, 2001).

The E1 reaction is a minimum two-step mechanism where the first step is the rate limiting step that involves loss of GSH from GSMOI to form a carbocation intermediate. In the second step a base removes the C-3 proton forming the double bond. The E2 reaction is a one-step concerted mechanism in which removal of the C-3 proton by a base and loss of GSH occur simultaneously. The E1 mechanism is highly unlikely due to the formation of an unstable primary carbocation intermediate, but the E1 and E2 mechanisms can be easily distinguished by studying the effect of pH on the reaction. (Smith, 2001).

The E1cb mechanism is also a two-step mechanism, where we have “fast” removal of the C-3 proton forming a stable conjugate base (from now onwards referred as cb) or carbanion intermediate. This is followed by a slow elimination step, forming GSH and MOI. The rate of an E1 mechanism is not expected to change with changes in pH as abstraction of the C-3 proton is not expected to be rate-limiting, while the rate of the E2 mechanism is predicted to have a proportional relationship with pH.



**Figure 7.** Possible elimination mechanisms for GSMOI, and predicted experimental observations, based on literature precedence.

The rates of both the E2 and E1cb reactions are expected to be pH dependent. However, the E1cb mechanism can be distinguished by the formation of the intermediate conjugate base. Formation of the cb can be observed by running the reaction in deuterated water (D<sub>2</sub>O) and analyzing kinetic data of proton exchange obtained from <sup>1</sup>H NMR spectrometry. For the two step E1cb mechanism, it is expected that fast proton exchange with D<sub>2</sub>O solvent will be observed by the disappearance of the C-3 proton <sup>1</sup>H NMR signal, but no (or much slower) GSH elimination and formation of MOI. For the one step E2 mechanism, the loss of the C-3 proton signal will occur at the same rate as the development of NMR signals for MOI.

#### ***1.4 Significance of this work***

The significance of this work can be directly related to the development of cancer therapeutic agents, an important area of study given that cancer is the cause of 1 in every 8 deaths worldwide, and research for new and different drugs is a top concern in drug development labs. The purpose of this research was to discover more relevant information about the inhibition of GxI by GSMOI that can be used in the future for drug development. If GSMOI is a mechanism-based inhibitor of GxI, then understanding how GSMOI undergoes elimination under physiological conditions is critical. By understanding this mechanism, predictions on how other enzymes might be affected by this inhibitor can be proposed. Overall this research will lead to a better understanding of the interactions of GSH, MOI, and GxI, and how these interactions can teach alternate approaches to one day curing cancer.

## ***2. Materials and Methods***

All reagents were purchased from Sigma-Aldrich or Fisher Scientific and used without further purification. Methanol was purchased at 99.9% purity. All glassware was washed with Micro-90 cleaner, and then rinsed with deionized water and acetone before oven drying at 80°C. Dry ice was purchased from Dry Ice Corp., Hingham, MA.

To study and characterize the elimination of GSH from GSMOI, kinetic assays were performed, in triplicate, in 50 mM potassium phosphate buffer at 25°C, using UV/Vis spectroscopy on a PerkinElmer Lambda 25 UV/vis spectrometer equipped with a PE PCB 1500 Water Peltier System for temperature control, and the UV WinLab software. To confirm the structure of the products, Nuclear Magnetic Resonance (<sup>1</sup>H NMR) was performed using a JEOL ECX-400M spectrometer. All NMR samples were prepared in 50 mM potassium phosphate buffer at a specified pH that had been exchanged into D<sub>2</sub>O solvent (pD = pH + 0.4).

### ***2.1 Synthesis of 3-Bromooxindole-3-acetic acid (BOAA)***

To prepare BOAA, 3.00 g of indole acetic acid, IAA, (17.1 mmol) were added to a 3-neck round bottom flask (RBF). The RBF was previously dried in the oven and covered with aluminum foil. To the same flask, 75ml of warm tert-butanol was added along with a magnetic stir bar. The IAA was stirred until dissolved, then 6.30 g of N-Bromosuccinimide (35.4 mmol) were added over a 60 minute time period. The color of the mixture changed from orange to reddish, and the mixture was stirred for an additional 10 minutes. The reaction was stored in the -80°C freezer overnight, and the next day was allowed to warm to room temperature before getting separated into different portions in a pear-shaped RBF. Each portion was put on the high vac in order to

evaporate the tert-butanol. After 30 minutes, all the tert-butanol was evaporated and the resulting syrup solution had a brown color. This was then dissolved in 50 mL of dry ethyl ether and magnesium sulfate was added as the drying agent. The mixture was stirred for 15 minutes and was filtrated to remove succinimide and the drying agent. The mixture was then placed on a rotavap for 20 minutes to evaporate the ether and dissolved again in 25 mL of ether. After 1 hour at room temperature, the mixture was filtrated again and concentrated on the rotavap. Residual ether entrapped in the brown syrup was evaporated on the high vac for approximately 10 minutes. The viscous syrup was dissolved in 25 mL of chloroform and stored for two hours at -20°C to allow for the precipitation of BOAA. The product was isolated by vacuum filtration with a Hirsch funnel. The product was scrapped onto a watch glass and put in a vacuum desiccator to dry overnight and then characterized by melting point and thin layer chromatography.

## ***2.2 Synthesis of 3-methyleoxindole (MOI) from 3-Bromooxindole-3-acetic acid (BOAA)***

To prepare MOI, 199 mg (0.736 mmoles) of BOAA, were dissolved in 1.5ml of 100% ethanol. This solution was then pipet filtrated (with cotton) to remove any insoluble material. The solution was transferred to a 10mL glass vial and stirred while cooling in an ice bath. While stirring, a 7mL ice cold solution containing 124 mg (1.48 mmoles) of sodium bicarbonate was added to the reaction mixture dropwise over a period of 3 minutes. The solution was allowed to stir for a couple of minutes until a yellow color was observed, and pH paper was used to verify that the solution was weakly acidic. The precipitate was filtrated using a Hirsch funnel, then quickly washed with 20mL of ice-cold deionized water to remove all traces of the sodium bicarbonate. The product was then transferred to a pre-weighed watch glass and placed in a vacuum

desiccator to dry overnight in the dark. The dried product was weighed, the percent yield was calculated, and the purity was determined using UV-Vis spectroscopy and  $^1\text{H}$  NMR spectrometry.

### ***2.3 Synthesis of Glutathione-3-Methyleneoxindole Conjugate (GSMOI) from 3-Methyloxindole and Glutathione.***

To make GSMOI, 140 ml of 25% methanol in deionized  $\text{H}_2\text{O}$  was added to a 250 ml Erlenmeyer flask. The solution was stirred under an atmosphere of argon for 15 minutes, then 70mg (0.228 mmole) of reduced glutathione (GSH) were added to the flask and stirred for an additional 5 minutes. The pH of the solution was adjusted using 1M potassium hydroxide to pH 6. Then, to the stirring solution, 39.7 mg (0.274 mmole) of 3-methyleneoxindole (MOI) were added in three portions over a period of 10 minutes. The solution was stirred for an additional one hour until all the yellow color from MOI dissipated. The reaction mixture was gravity filtrated into a clean flask to remove any white precipitate due to MOI polymerization, and the filtrate was cooled in ice. The filtrate was concentrated to about 20ml on the high vac at  $30^\circ\text{C}$  to remove methanol, and the concentrated aqueous GSMOI solution was then lyophilized to dryness.



**Figure 8.** Lyophilizer used to dry the GSMOI.

The product was obtained as a fluffy, white powder which was then washed with anhydrous ether two times to extract excess MOI. The ether was removed with a pipet and the white solid dried under a gentle stream of dry argon for 5 minutes. The product was transferred to a pre-weighed watch glass and dried thoroughly in a vacuum desiccator overnight in the dark. Then GSMOI was stored under dry conditions at -20°C. The yield of the dry product was determined, and the structure verified by UV-Vis spectroscopy and <sup>1</sup>H NMR spectrometry.

#### ***2.4 Characterization using NMR***

<sup>1</sup>H NMR was used to determine the purity of both MOI and GSMOI. For MOI 6 mg of solid sample was dissolved in Acetone d<sub>6</sub>. GSMOI NMR samples were prepared in two ways. Method I: 7-10 mg of GSMOI were dissolved in 1ml of D<sub>2</sub>O and lyophilized to dryness. This process was repeated two more time with fresh D<sub>2</sub>O to exchange acidic H for D. Then the dry, exchanged sample was dissolved in 1 ml of Deuterated Potassium Phosphate, (D)KP buffer at a specified pD. Method II: 7-10 mg of GSMOI were dissolved directly in 1ml of (D)KP buffer at a specified pD. The (D)KP was prepared by exchanging 10 mL of (H)KP buffer at a specified pH with D<sub>2</sub>O on the high vac. Samples were flushed with dry argon to limit exposure to air. The pD correction factor is +0.4 units, so for a pH 7 buffer, the pD is 7.4.

#### ***2.5 Development of UV-VIS Spectroscopic Methods for Elimination Kinetics Studies***

A stock solution of GSMOI was prepared in deionized (DI) water (4.86 mM). The stock solution was kept on ice during use, and stored at -20°C. Under these conditions this solution was

stable for at least 4 weeks. A 5 mM stock solution of 5,5-dithio-bis-(2-nitrobenzoic acid) (DTNB, Ellman's reagent) was prepared in KP buffer, kept on ice during use, and stored at 4°C.

Method I. 50 mM KP buffer in a quartz cuvette (1.0 mL) was pre-incubated for 10 minutes at 25°C. To study the rate of GSH elimination, the reaction was initiated by adding an aliquot of the GSMOI stock solution to a final concentration of 50 µM. Repeat wavelength scans (200-500 nm) were done at specified time intervals for 3 hours. To determine the rate of formation of MOI the change in absorbance at 248 nm was analyzed. Due to the low extinction coefficient of MOI at 248 nm this method did not provide the sensitivity for kinetic analysis. We switched to Elman's Reagent, DTNB (5,5-dithio-bis-(2-nitrobenzoic acid) as described below.

Method II. A 5 mM solution of DTNB was pre-incubated with 50 mM KP buffer in a quartz cuvette at 25 °C. After equilibration for 5 minutes, reaction was initiated by the addition of various concentrations of the GSMOI stock solution. Initial rate slopes from absorbance (412 nm) vs time curves were determined over the initial 10-15% of reaction. The extinction coefficient of DTNB at 412 nm is (14150 m<sup>-1</sup> cm<sup>-1</sup>).

## ***2.6 Determination of Reaction Order***

To determine the order of the reaction, a 5 mM solution of DTNB was incubated with 50 mM potassium phosphate buffer in a quartz cuvette at 25°C. After temperature equilibration for at least 5 minutes, the reaction was initiated by the addition of different aliquots of the GSMOI stock solution (10, 20, 40 and 80 µL). Absorbance vs time curves at 412 nm were recorded and initial rate slopes from the first 10-15% of the reaction were used to calculate reaction rates.



## ***2.7 Effect of pH on the rate of GSH elimination***

To determine the effect of pH on the reaction, a 5mM solution of DTNB was incubated with 50 mM potassium phosphate buffer (at pH 6.50, 7.00 and 7.50) in a quartz cuvette at 25°C. After equilibration for 5 minutes, the reaction was initiated by the addition of different aliquots of GSMOI stock solution to the cuvette. The change in absorbance at 412 nm was recorded and initial 10-15% of the reaction was used in order to collect the initial slopes from absorbance vs time curves.

## ***2.8 Effect of potential catalysts***

The DTNB assay used in section 2.6, was used again in order to determine the effect of different bio-molecules as potential catalysts for the elimination reaction. In addition to DTNB, a 250 mM solution of the potential catalyst was incubated with 50 mM potassium phosphate buffer in a quartz cuvette at 25°C. After equilibration for 5 minutes, the reaction was initiated by the addition of various concentrations of GSMOI. The absorbance at 412 nm was recorded and initial 10-15% of the reaction was used in order to collect the initial slopes from absorbance vs time curves.

## ***2.9 Effect of buffer concentration on rate constant $k$***

A 5 mM solution of DTNB was incubated with 10, 50 and 250 mM potassium phosphate buffer in a quartz cuvette at 25 °C (pH 7). After equilibration for 5 minutes, reaction was initiated by the addition of various concentrations of GSMOI to the cuvette. Initial rate slopes from absorbance vs time curves were determined over the initial 10-15% of reaction.

### ***2.10 GSMOI C-3 Proton exchange kinetics***


All glassware was dried in an oven at 100°C, and stored in a desiccator. Dry argon was used to flush all samples and protect from air. 7-10 mg of GSMOI were dissolved in 1ml of (D)KP buffer at pD 6.69 and 7.34. A stopwatch was started upon addition of D-KP buffer to the GSMOI samples, and these solutions were immediately transferred to an NMR tube and scans started within 2-3 minutes. The scans were collected every 8 minutes for approximately one hour, and the C-3-H and cys- $\alpha$ -H signals were integrated.

### 3. Results and Discussion

#### 3.1 Synthesis of 3-Bromooxindole-3-acetic acid (BOAA) and 3-methyloxindole (MOI)

Because access to an NMR was restricted at the time, BOAA was only characterized by observing the melting point range of 153-158°C. This compares very well with the literature mp of 158°C.


After drying in a vacuum desiccator overnight, the MOI sample looked bright yellow, and the % yield was 42.5%. The <sup>1</sup>H NMR in Figure 9 confirmed no impurities, as all the peaks were consistent with the structure of MOI. The UV-Vis showed a characteristic doublet absorption band at 248 and 252 nm, characteristic with MOI, Figure 13.




**Figure 9.** <sup>1</sup>H NMR spectrum of 6 mg of MOI in Acetone d<sub>6</sub>.

### 3.2 Synthesis of Glutathione-3-Methyleneoxindole Conjugate (GSMOI) from 3-Methyloxindole and Glutathione

GSMOI was obtained as a dry white powder which was kept under dry conditions in the freezer, and the UV-Vis spectrometer confirmed that there was not MOI residue in the sample. Although GSH is a tripeptide, analysis by  $^1\text{H}$  NMR is straight forward as seen in the  $^1\text{H}$  NMR of 7 mg of GSH in 50 mM (D)KP buffer, pD 7.34, Figure 10. The  $^1\text{H}$  NMR for GSMOI showed no presence of MOI, as all the peaks were consistent with the GSMOI structure, Figure 11. It's interesting that the C-3 oxindole proton is not observed at 3.8 ppm, suggesting that it has undergone fast exchange with the  $\text{D}_2\text{O}$  solvent at pD 7.34.



**Figure 10.**  $^1\text{H}$  NMR spectrum of 7 mg of GSH in 1 mL of 50 mM (D)KP buffer, pD 7.34.




**Figure 11.**  $^1\text{H}$  NMR spectrum of 7 mg of GSMOI in 1 mL of 50 mM (D)KP buffer, pD 7.34

### ***3.3 Development of UV-VIS Spectroscopic Methods for Elimination Kinetics Studies***

The elimination reaction of GSMOI produces two products, GSH and MOI, Figure 7. Two methods were evaluated to obtain reproducible kinetic data for the elimination of GSH from GSMOI. In Method I we attempted to monitor the formation of MOI based on its absorption band at 248 nm, see Figure 13. Although Method I did provide us with preliminary data, reproducibility was a problem. As MOI has a strong extinction coefficient at 248 nm ( $\epsilon_{248} = 29,000 \text{ M}^{-1}\text{cm}^{-1}$ ), the problem may be the reactivity of free MOI toward Michael addition reactions with nucleophilic compounds, and its polymerization at neutral to basic conditions.

Method II used 5,5-dithio-bis-(2-nitrobenzoic acid) (Ellman's Reagent, DTNB) which is a sensitive reagent to detect and quantify free thiols based on thiol-disulfide exchange reaction, Figure 12. The  $\text{TNB}^-$  anion has a visible yellow color and strong extinction coefficient at 412 nm

( $\epsilon_{412} = 14,150 \text{ M}^{-1}\text{cm}^{-1}$ ). Method II appeared to give reproducible data. There was also no evidence that the  $\text{TNB}^-$  anion was reacting with the free MOI by Michael addition. At the end of the reaction with DTNB, the absorbance at 412 nm was stable, suggesting little reaction with the free MOI. The  $\text{TNB}^-$  thiolate anion is expected to be a poor nucleophile due to resonance stabilization of the thiolate anion with the electron withdrawing nitro group.




**Figure 12.** Thiol-disulfide exchange reaction between glutathione and DTNB producing the  $\text{TNB}^-$  ion to assay for the formation of GSH in the elimination reaction of GSMOI.

### 3.4 UV-VIS titration of MOI with GSH

This experiment was done when we did not have access to an NMR, in order to evaluate the purity of the MOI being used to synthesize GSMOI. The UV-Vis titration of  $50 \mu\text{M}$  MOI was conducted in  $50 \text{ mM}$  KP buffer at pH 7. This sample was titrated by the addition of approximately  $10 \mu\text{M}$  increments of GSH, and the UV-Vis scans taken from 200-400 nm as shown in Figure 13.

The “doublet” band for MOI at 248 & 250 nm is seen to clearly decrease with each addition of MOI. No additional change occurred upon GSH concentrations above 49  $\mu\text{M}$ , suggesting complete titration of the MOI, and an estimated purity of 95%. In addition, this experiment demonstrates that the formation of GSMOI from addition of GSH to MOI occurs without any complications from side products. With incremental addition of GSH we observed a smooth decrease in the absorption band of MOI, plus the observation of at least three, clear isobestic points.




**Figure 13.** UV-Vis titration of 50.2  $\mu\text{M}$  MOI with approximately 10  $\mu\text{M}$  increments of GSH in 50 mM KP buffer, pH 7.

### 3.5 Reaction order

Preliminary results using the MOI UV-Vis assay suggested that the elimination reaction occurred by a mixed order mechanism as we could not distinguish between zero, 1<sup>st</sup> or 2<sup>nd</sup> order kinetics. It is unusual for elimination reactions to be mixed order, and typically the elimination reaction is observed under pseudo first order conditions with limiting GSMOI. We attributed this

preliminary result to problems with the UV-Vis assay where we were observing the decrease in the MOI absorbance band at 248 nm.

Using the DTNB UV-Vis assay to monitor the elimination of GSH produced more consistent data. In Figure 14, we see that by increasing the concentration of GSMOI from 0.50 to 4.0 mM, that the initial rate for elimination increases in a corresponding manner, suggesting that zero order kinetics are not being observed. A plot of natural log of initial rate vs time gave a straight line which is consistent with pseudo first order kinetics.



**Figure 14.** UV-Vis assay using DTNB to monitor the elimination of GSH from GSMOI using 0.50 – 4.0 mM GSMOI in 50 mM KP buffer, pH 7.

### 3.6 Effect of pH

The effect of pH on the rate of GSH elimination was an important experiment in distinguishing between the three possible mechanisms shown in Figure 7 above. In an E1 carbocation mechanism, proton transfer follows rate limiting elimination of the GSH leaving



group, so pH should have little effect on rate. The concerted E2 mechanism requires proton transfer at the same time as GSH elimination, so the reaction rate would be expected to increase with increasing pH. The rate of an E1cb reaction is also expected to increase with increasing pH as the fast proton transfer step is followed by slow GSH elimination.

A 5mM solution of DTNB was incubated in 50 mM potassium phosphate buffer (at pH 6.50, 7.00 and 7.50) in quartz cuvettes at 25°C. The reaction was initiated by the addition of various concentrations of GSMOI to the cuvette. The change in absorbance at 412 nm was recorded and the initial 10-15% of the reaction was used in order to collect the initial slopes from absorbance vs time curves. The pseudo 1<sup>st</sup> order rate constants with standard deviations at each pH are presented in Table I, and indicate that the rate constants increase with increasing pH. This suggests that changes in pH affect the rate of reaction, and that elimination is following either an E2 or E1cb mechanism. In our preliminary experiments using the less sensitive MOI assay at 248 nm, we observed little change in elimination reaction rate between pH 6.50-7.50, but a significant decrease in the reaction rate at pH 6.00. We believe that this may suggest a change in the rate limiting step consistent with an E1cb elimination mechanism because the “fast” proton transfer step is now competing with GSH elimination. Although the DNTB assay is more reproducible, and the data follow a similar trend, the decrease in rate with decreasing pH was not as pronounced. We believe that this discrepancy results from the DTNB assay being less sensitive below pH 7, as protonation of the TNB<sup>-</sup> thiolate ion inhibits conjugation and decreases production of the yellow color at 412 nm. The data presented in this table are reliable because the standard deviations were all close to  $\pm 10\%$  suggesting the rate effects are not due to experimental error.

**Table I.** Variation of pseudo first order rate constants for the elimination reaction of GSH from GSMOI using the DTNB assay with 50 mM KP buffer at different pH values.

<b>pH</b>	<b>Rate constant k (1/sec) x 10<sup>-3</sup></b>	<b>Standard deviation x 10<sup>-3</sup></b>
6.5	36.5	4.20
7.0	50.5	1.36
7.5	64.3	5.06

### ***3.7 Effect of potential catalysts***

Our goal is to understand the chemical mechanism for GSH elimination from GSMOI, as a potential model for the elimination being catalyzed under physiological conditions. This led us to examine the effect of bio-molecules as potential catalysts for the elimination reaction. Our standard kinetic assay had 5 mM DTNB incubated in 50 mM potassium phosphate buffer at pH 7 in a quartz cuvette at 25°C. The cuvette also included a large excess (250 mM) of the potential catalyst to preserve pseudo 1<sup>st</sup> order conditions. The reaction was initiated by the addition of various concentration of GSMOI to the cuvette. The change in absorbance at 412 nm was recorded and initial 10-15% of the reaction was used in order to collect the initial slopes from absorbance vs time curves. The pseudo 1<sup>st</sup> order rate constants with standard deviations for each catalyst are presented in Table II, along with a control containing no catalyst. The data presented in this table is reliable because the standard deviations were all close to  $\pm 10\%$  suggesting the effects are not due to experimental error.

**Table II.** Variation of pseudo 1<sup>st</sup> order rate constants after incubation GSMOI with various biomolecules as potential elimination catalysts.

<b>Catalyst</b>	<b>Rate constant k (1/sec) x 10<sup>-3</sup></b>	<b>Standard Deviation x 10<sup>-3</sup></b>
L-Histidine	25.6	3.38
Glycine	24.7	5.00
Guanidine	23.0	6.77
<b>Control</b>	<b>50.5</b>	<b>1.36</b>

Compared to the control reaction with no “catalyst”, the data in Table II indicate that these potential catalysts all actually decrease the reaction rate constant. This result was a little surprising. These molecules, acting as bases, might have been expected to either increase the rate of an E2 mechanism, or have little to no effect on an E1cb mechanism. The inhibition is interesting, and may suggest that these additives are affecting the DTNB reaction with free GSH. Another possible explanation is that the GSH portion of GSMOI is involved in intramolecular catalysis of the elimination, which is being inhibited through ion pairing with the bases in Table II. More research is required on this observation to better understand these results.

### ***3.8 Effect of buffer concentration on rate constant $k$***

If the GSH portion of GSMOI is involved in intramolecular catalysis of GSH elimination, changing the buffer concentration may have an effect on the reaction rate. Increasing the buffer concentration will increase ionic strength, resulting in ion pairing with GSH and reaction inhibition. On the other hand, decreasing the buffer concentration may potentially increase the effect of GSH catalysis. Various concentrations of GSMOI were incubated with 10, 50 and 250 mM KP buffer at pH 7. The pseudo 1<sup>st</sup> order rate constants and standard deviations are shown in Table III.

**Table III.** Effect of KP buffer concentration on the rate of GSMOI elimination reaction, pH 7.

<b>Buffer Concentration</b>	<b>Rate constant <math>k</math> (1/sec) x 10<sup>-3</sup></b>	<b>Standard Deviation x 10<sup>-3</sup></b>
10 mM	6.27	0.509
50 mM	50.5	1.36
250 mM	23.4	4.55

As compared to the standard reaction conditions with 50 mM KP buffer, either increasing or decreasing the buffer concentrations resulted in an inhibition of the elimination reaction. Inhibition at higher buffer concentration is understandable due to ion pairing with GSH, but


inhibition at lower buffer concentration is confusing. KP buffers are not known to act as catalysts, and it is unlikely that changing the buffer concentrations are effecting the DTNB assay. Maybe at low concentrations of buffer, there is intermolecular ion-pairing of the GSH parts of GSMOI pairs, and this might be expected to inhibit GSH catalysis. Again, this is an interesting result, but additional experimentation work is needed to understand these data. The data presented in this table are reliable because the standard deviations were all close to  $\pm 10\%$  suggesting the effects are not due to experimental error.

### ***3.9 GSMOI C-3 Proton exchange kinetics***

The most direct way to distinguish between the E2 and E1cb mechanisms was to attempt to observe removal or exchange, respectively, of the C-3 GSMOI oxindole proton using  $^1\text{H}$  NMR in (D)KP buffer. For an E2 mechanism, as the C-3 proton is removed and its  $^1\text{H}$  NMR signal decreases, there should be a corresponding increase in the vinyl protons of MOI at 6.2 ppm. On the other hand, an E1cb mechanism will result in a reversible, fast exchange of the C-3 proton with deuterium, its  $^1\text{H}$  NMR signal will quickly decrease, but with very little MOI being produced.


A 7-mg sample of GSMOI was dissolved in 1ml of (D)KP buffer at pD 6.69. The  $^1\text{H}$  NMR spectrum is shown in Figure 15. As compared to the GSMOI  $^1\text{H}$  NMR spectrum obtained at pD 7.34 where the C-3 proton had rapidly exchanged and was not observed (Figure 11), at pD 6.69 the C-3 oxindole proton is clearly visible at 3.8 ppm. Integration of the C-3 proton (b) was compared to the integration of the cys- $\alpha$ -H signal (a) at 4.25 ppm, and it was found that within 2 minutes of initiating the reaction that the C-3 signal had decreased by 31%, but only a trace of MOI was observed based on the vinyl proton signal at 6.8 ppm. A stopwatch was started upon

addition of D-KP buffer to the GSMOI sample, and NMR scans started within 2-3 minutes. Scans were collected every 8 minutes for approximately one hour, and the C-3-H and cys- $\alpha$ -H signals were observed.



**Figure 15.** <sup>1</sup>H NMR spectrum of 7 mg of GSMOI in 1 mL of 50 mM (D)KP buffer, pD 6.69.

The partial NMR spectra in Figure 16 show only the C-3-H signal at 3.8 ppm and cys- $\alpha$ -H signal at 4.25 ppm. Over a period of 26 minutes, the C-3 oxindole proton was observed to undergo time-dependent exchange with D<sub>2</sub>O solvent. This exchange is much faster than GSH elimination as only a trace of MOI was observed over this same time period (data not shown). ***This result, in combination with the data above, is strong evidence that that chemical mechanism for the elimination of GSH from GSMOI follows an E1cb mechanism.***



**Figure 16.** <sup>1</sup>H NMR spectrum of 7 mg of GSMOI in 1 mL of 50 mM (D)KP buffer, pD 6.69 undergoing time dependent exchange of the oxindole C-3 proton.

## ***4. Conclusions***

The significance of this work can be directly related to the development of cancer therapeutic agents, an important area of study given that cancer is the cause of 1 in every 8 deaths worldwide, and research for new and different drugs is a top concern in drug development labs. The purpose of this research is to discover more relevant information about the inhibition of GxI by GSMOI that can be used in the future for drug development.

Previous students in Dr. Brush's research group found preliminary evidence that GSMOI is both a competitive and mechanism-based inhibitor of GxI, however more study is needed to better understand those findings. MOI is not a good choice as a therapeutic agent because of its poor aqueous solubility, and non-selectivity as an alkylating agent. By conjugating MOI with GSH to form GSMOI, we believe this compound will specifically target GxI and inhibit by mechanism-based inactivation.

Once bound to the active site, we think that GxI will catalyze the elimination of GSH from GSMOI, releasing MOI. As MOI is an alkylating agent for cysteine, we believe that the MOI will selectively alkylate an active site cysteine residue by Michael addition, Figure 6 (above). The release of MOI and GSH from GSMOI is an example of an elimination reaction. There is a leaving group (GSH), loss of a proton and the formation of a product with a double bond (MOI). If GSMOI is a mechanism-based inhibitor of GxI, we needed to understand how GSMOI undergoes elimination under physiological conditions.

The main outcome of this thesis project is that we now have a better understanding of the mechanism by which GSH is eliminated from GSMOI. We developed an assay to quantitatively study the rate of the elimination reaction, under conditions that would allow us to study factors that affect the rate limiting step. We have studied how the reaction rate was affected by pH and potential catalysts. Our rate data allowed us to distinguish between the three most common biological elimination mechanisms that could explain the elimination of GSH and formation of MOI: E1, E2 or E1cb, Figure 7. Our key data are summarized in Table IV. The E1 reaction is a minimum two-step mechanism where the first step is the limiting rate step where the GSH leaving group leaves, forming a carbocation intermediate. In the second step a base removes the C-3 proton forming the double bond. The E1 mechanism is highly unlikely due to the formation of an unstable primary carbocation intermediate.

**Table IV.** Summary of the outcome of the most significant experiments. The data strongly suggests E1cb mechanism.


<b>Experiment</b>	<b>Observed Results</b>	<b>Mechanism</b>
Reaction	1 <sup>st</sup> and 2 <sup>nd</sup>	E2 and E1cb
pH	Faster with higher pH	E2 and E1cb
Catalyst	Inhibition	Inconclusive
C-3 proton exchange	Fast exchange	E1cb

The E2 reaction is a one-step concerted mechanism. The base removes the C-3 proton and the GSH leaving group leaves at the same time forming the MOI double bond. The E1 and E2 mechanisms were distinguished based on the observation of elimination rate increase with increasing pH. The rate of an E1 mechanism was not expected to change with changes in pH, while the rate of the E2 mechanism was predicted to have a proportional relationship with pH.



The E1cb mechanism is also a two-step mechanism, where we have fast removal of the C-3 proton forming a stable conjugate base “cb” intermediate. This is followed by a slow elimination step, forming GSH and MOI. The rates of both the E2 and E1cb reactions were expected to be pH dependent. However, the E1cb mechanism was identified based on data suggesting formation of the “cb” intermediate based on the observation of fast proton exchange in D<sub>2</sub>O solvent by <sup>1</sup>H NMR. We observed the disappearance of the C-3 proton <sup>1</sup>H NMR signal, and much slower formation of MOI, Figure 16. This experimental result suggests against the one step E2 mechanism, where the loss of the C-3 proton signal should have occurred at the same rate as the formation of MOI.


The goals of this thesis project were achieved as the data suggest that the elimination mechanism is E1cb. What do the results of this research suggest for GSMOI as a mechanism based inhibitor of GxI? In its “normal” reaction GxI catalyzes the conversion of the thiohemiacetal of GSH and MG into (S)-D-lactoylglutathione, Figure 2 (above). The chemical mechanism involves a proton transfer with an intermediate carbanion, as shown in Figure 17 (Chari and Kozarich, 1981).



**Figure 17.** The proton transfer chemical mechanism of GxI converting the GSH-MG hemiacetal substrate into lactoylglutathione.

The GxI proton transfer mechanism with its normal substrate suggests that GxI may also catalyze a proton transfer for the E1cb elimination of GSH from GSMOI. MOI is also produced, which we believe may then alkylate an active site cysteine and lead to irreversible inactivation. This means that GSMOI would also be a substrate, or mechanism-based inhibitor for GxI. This idea may work as human GxI has been found to catalyze proton transfer reactions for unusual substrates such as those shown in Figure 18 (Chari and Kozarich, 1981, 1983; Landro, Brush and Kozarich, 1992). Its also exciting that the fluoromethylglyoxal GSH-thiohemiacetal undergoes E1cb elimination of fluoride ion, catalyzed by human GxI (Chari and Kozarich, 1981). This may suggest that human GxI is capable of catalyzing GSH elimination from GSMOI!

Now armed with a better understanding of the aqueous chemistry of GSMOI, we will be able to better predict if other enzymes might also be affected by this molecule. Overall this research will lead to a better understanding of the interactions of GSH, MOI, and GxI and how these interactions can teach us alternate approaches to one day curing cancer.



**Figure 18.** Alternative substrates for human GxI.

## ***5. Future Work***

As to any research project, there are still a few things that need to be done in the future. The effect of buffer concentration changes were surprising, with various effect on the rate of reaction. At lower buffer concentration, the reaction rate is very small, and it increases as the buffer strength is increased. However, at a much higher concentration, this increasing pattern is lost. Therefore, more studies need to be done in order to determine if the buffer concentration is affecting the DTNB reaction rather than the elimination mechanism.

Another assumption made while conducting this research was that TNB anion does not react with MOI, and this needs to be verified experimentally.

Also, the addition of catalyst to the reaction was expected to have little to no effect on the reaction rate since it is an E1cb reaction. The first step, or removal of proton is very fast while the second step, elimination of leaving group is very slow. The addition of catalyst appears to be inhibiting the reaction rate, and therefore more work needs to be done in order to understand the reasons behind it. A future experiment that can be done to complete this study is analyze the role of GSH backbone in elimination mechanism. This would involve the synthesis of a smaller MOI conjugates, such as N-acetyl-cysteine-MOI and N-acetyl-cysteamine-MOI.

## 6. Bibliography

Bair III, Warner B., Christopher M. Cabello, Koji Uchida, Alexandra S. Bause, and Georg T. Wandrak. "GLO1 Overexpression in Human Malignant Melanoma." *Melanoma Research* 20.2 (2010): 85-96. *NCBI*. Web. 27 Feb. 2017. <<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2891514/pdf/nihms208136.pdf>>.

Brush, E., Goldberg J., Petrounia I. (1994). "Transient Inactivation of Almond Mandelonitrile Lyase by 3-Methyleneoxindole." *Biochemistry*. 33: 2891-99

Chauhan, Swati C., and Rental Madhubala. "Glyoxalase I Gene Deletion Mutants of Leishmania Donovanii Exhibit Reduced Methylglyoxal Detoxification." *PLoS One* (2009): n. pag. *NCBI*. Web. 27 Feb. 2017. <<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2728510/pdf/pone.0006805.pdf>>.

Guo, Yi, Yuning Zhang, Xunjun Yang, Panpan Lu, Xijuan Yan, Fanglan Xiao, Huaibin Zhou, Chaowei Wen, Mengru Shi, Jianxin Lu, and Qing H. Meng. "Effects of Methylglyoxal and Glyoxalase I Inhibition on Breast Cancer Cells Proliferation, Invasion, and Apoptosis through Modulation of MAPKs, MMP9, and Bcl-2." *Cancer Biology & Therapy* 17.2 (2016): 169-80. *NCBI*. Web. 27 Feb. 2017. <<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4848000/pdf/kcibt-17-02-1121346.pdf>>.

Igor Allaman, Mireille Bélanger and Pierre J. Magistretti (2015), "Methylglyoxal, the dark side of glycolysis", *Front. Neurosci.* 9 (23): 1-12. Web. 25 April 2018. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4321437/pdf/fnins-09-00023.pdf>

Landro, JA, Brush, EJ, and Kozarich, JW (1992), "Isomerization of (R)- and (S)- Glutathiolactaldehydes by Glyoxalase I: The Case for Dichotomous Stereochemical Behavior in a Single Active Site?", *Biochemistry*, 131, 6069-6077.

Jaetaek Kim, Jang-Won Son, [...], and Soon-Hyun Shinn. "Methylglyoxal Induces Apoptosis Mediated by Reactive Oxygen Species in Bovine Retinal Pericytes."(2001)

Okado A, Kawasaki Y, Hasuike Y, Takahashi M, Teshima T, Fujii J, Taniguchi N." Induction of apoptotic cell death by methylglyoxal and 3-deoxyglucosone in macrophage-derived cell lines." *Biochem Biophys Res Commun.* 1996;225:219-224. [[PubMed](#)]

Osborne BA. Apoptosis and the maintenance of homeostasis in the immune system. *Curr Opin Immunol.* 1996;8:245-54. [[PubMed](#)]

Ravi V. J. Chari and John W. Kozarich (1981), "Deuterium Isotope Effects on the Product Partitioning of Fluoromethylglyoxal by Glyoxalase I", *J. Biol. Chem.*, 256 (19): 9785-9788.

Ravi V. J. Chari and John W. Kozarich (1983), "Glutathiohydroxyacetone: 1H NMR Determination of the Stereochemistry of Proton Exchange by Glyoxalase I. Evidence for a cis Eneiol Intermediate Based on Mirror-Image Catalysis", *J. Amer. Chem. Soc.*, 105, 7169-7171.

Smith, B. Michael, March Jerry, "Advanced Organic Chemistry, Reactions, Mechanisms and Structures." John Wiley & Sons, Inc 2001.

Thornalley, Paul J. "The Glyoxalase System: New Developments towards Functional Characterization of a Metabolic Pathway Fundamental to Biological Life\*." *Biochemistry Journal* 269 (1990): 1-11. *NCBI*. Web. 27 Feb. 2017.  
<<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1131522/pdf/biochemj00180-0011.pdf>>.

Vander Jagt DL, Hunsaker LA, Vander Jagt TJ, Gomez MS, Gonzales DM, Deck LM, Royer RE. "Inactivation of glutathione reductase by 4-hydroxynonenal and other endogenous aldehydes". *Biochem Pharmacol.* 1997;53:1133–1140. [[PubMed](#)]

White MK, Cinti C. A morphologic approach to detect apoptosis based on electron microscopy. *Methods Mol Biol.* 2004; 285:105–11. [[PubMed](#)]

Wu L, Juurlink BH." Increased methylglyoxal and oxidative stress in hypertensive rat vascular smooth muscle cells." *Hypertension.* 2002;39:809–814. [[PubMed](#)]